Contents Volume 11, 1986/1987

Introduction
P. De Wever (Paris, France) 1
Trends in the development of Paleozoic Radiolaria
B.B. Nazarov (Moscow, U.S.S.R.) and A.R. Ormiston (Tulsa, OK, U.S.A.) 3
Origin and biostratigraphic potential of the stauraxon polycystine Radiolaria
B.B. Nazarov (Moscow, U.S.S.R.) and A.R. Ormiston (Tulsa, OK, U.S.A.) 33
Some Late Permian radiolarians from pelitic rocks of the Tatsuno Formation (Hyogo Prefecture), southwest Japan
M. Caridroit (Osaka, Japan) and P. De Wever (Paris, France) 55
A newly proposed radiolarian zonation for the Jurassic of Japan
A. Matsuoka and A. Yao (Osaka, Japan) 91
New data on the biostratigraphy of the Tuscan Cherts at Monte Cetona (Southern Tuscany, Italy)
M. Conti (Florence, Italy) 107
Datation par les radiolaires de la Formation des Radiolarites s.s. de la série du Pinde—Olonos (Grèce): Bajocien (?)—Tithonique. [Radiolarian dating of the Radiolarite Formation s.s. of the Pindos—Olonos Zone (Greece): Bajocian (?)—Tithonian]
P. De Wever and F. Cordey (Paris, France) 113
The onset of radiolarian deposition in the ophiolite sequences of the Northern Apennines
M. Conti (Florence, Italy) and M. Marcucci (Florence, Italy) 129
Middle to Late Cretaceous radiolarian zonation of the Bering region, U.S.S.R.
V.S. Vishnevskaya (Moscow, U.S.S.R.) 139
Morphological characters for a natural classification of Cenozoic Radiolaria, reflecting phylogenies
W.R. Riedel and A. Sanfilippo (La Jolla, CA, U.S.A.) 151
Final stages of skeletogenesis and early stages of disintegration for modern polycystine radiolaria
K.R. Bjørklund (Bergen, Norway) and R.M. Goll (Broken Arrow, OK, U.S.A.) 171
Evolution of the *Antarctissa* group
M.G. Petrushevska (Leningrad, U.S.S.R.) 185
Transportation of radiolarian shells by currents (calculations based on the example of the Kuroshio)
I.M. Popova (Vladivostok, U.S.S.R.) 197
A comparative analysis of the morphogenesis and morphometric diversity of mature skeletons of living *Didymocyrtis tetrathalamus tetrathalamus* and *Hexalonche amphisiphon*
O.R. Anderson (Palisades, NY, U.S.A.), C. Hemleben (Tübingen, F.R.G.), M. Spindler (Bremerhaven, F.R.G.) and J.L. Lindsey (Palisades, N.J., U.S.A.) 203
A refined radiolarian biostratigraphy for the Pleistocene of the temperate Indian Ocean
J.P. Caulet (Paris, France) 217
The radiolarian fauna of western Norwegian fjords: patterns of abundance in the plankton
N.R. Swanberg (Palisades, NY, U.S.A.) and K.R. Bjørklund (Bergen, Norway) 231
Size and shape patterns in the Recent radiolarian genus *Antarctissa* from a south Indian Ocean transect
A. Granlund (Stockholm, Sweden) 243

Differential dissolution of Upper Cretaceous planktonic foraminifera from a temperate region of the South Atlantic Ocean
B.A. Malmgren (Uppsala, Sweden) 251
Postglacial environmental change of the Pacific Ocean off the coast of central Japan
K. Chinzei (Kyoto, Japan), K. Fujioka (Tokyo, Japan), H. Kitazato (Shizuoka, Japan), I. Koizumi (Toyonaka, Japan), T. Oba (Kanazawa, Japan), M. Oda (Kumamoto, Japan), H. Okada (Yamagata, Japan), T. Sakai (Utsunomiya, Japan) and Y. Tanimura (Tokyo, Japan) 273
Valanginian to Aptian calcareous nannofossil stratigraphy and correlation with the upper M-sequence magnetic anomalies
T.J. Bralower (La Jolla, CA, U.S.A.) 293
Sedimentation of biogenic siliceous particles in Antarctic waters from the Atlantic sector
R. Gersonde (Bremerhaven, F.R. Germany) and G. Wefer (Bremen, F.R. Germany) 311

The pre-cephalic development of the skeleton of *Amphimelissa setosa* (Actinopoda: Nassellarida)
N.R. Swanberg (Palisades, NY, U.S.A.) and K.R. Bjørklund (Oslo, Norway) 333
The distribution of recent benthic foraminifera in the Polar Front region, southwest Atlantic
G..A. Mead (La Jolla, CA, U.S.A.) and J.P. Kennett (Kingston, RI, U.S.A.) 343

Errata 361